

* Chp. 5 Eigen-stuff

Def: When $T: V \rightarrow V$ is a lin. trans.
and we find a specific vector
 $\vec{x} \in V$ such that $\vec{x} \neq \vec{0}$

and $T(\vec{x}) = c\vec{x}$ for some
constant c

then we call \vec{x} an **eigenvector**
with **eigenvalue** c (often use $c = \lambda$,
(λ can be 0, but $\vec{x} \neq \vec{0}$)

(if T is just multiplying every
vector by a constant, then every
vector in V is an eigenvector, with
that constant λ its eigenvalue.)

However, most lin. trans. $T: V \rightarrow V$ have only
certain eigenvectors and eigenvalues. Find them!

Steps:

1) We work with $A = [T]_{\mathcal{B}}$

2) Let $A\vec{x} = \lambda\vec{x}$ ($\vec{x} \neq \vec{0}$)
then $\Rightarrow A\vec{x} = (\lambda I)\vec{x}$ ($\lambda I = \begin{bmatrix} \lambda & 0 & 0 & \dots & 0 \\ 0 & \lambda & & & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & 0 & 0 & \dots & \lambda \end{bmatrix}$)
 $\Rightarrow A\vec{x} - (\lambda I)\vec{x} = \vec{0}$
 $\Rightarrow (A - \lambda I)\vec{x} = \vec{0}$

So $\vec{x} \neq \vec{0}$ and $\vec{x} \in N(A - \lambda I)$

$\Rightarrow \det(A - \lambda I) = 0$

3) this gives us an algebraic equation
to solve for λ . Then plug back in to find \vec{x} .

ex) Let $T: p^2 \rightarrow p^2$

be given by $T(f(x)) = 2xf'(x) + 3xf''(x)$

Find the eigenvalues and their corresponding eigenvectors for T .

$\vec{e}_i \in \mathcal{E}$	$f'(x)$	$f''(x)$	$T(\vec{e}_i)$
1	0	0	0
x	1	0	$2x$
x^2	$2x$	2	$4x^2 + 6x$

$$1) \quad A = [T]_{\mathcal{E}}^{\mathcal{E}} = \begin{bmatrix} [0]_{\mathcal{E}} & [2x]_{\mathcal{E}} & [4x^2+6x]_{\mathcal{E}} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 2 & 6 \\ 0 & 0 & 4 \end{bmatrix}$$

$$2) \quad \det(A - \lambda I) = \det \left(\begin{bmatrix} 0 & 0 & 0 \\ 0 & 2 & 6 \\ 0 & 0 & 4 \end{bmatrix} - \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} \right) = 0$$

$$= \det \begin{pmatrix} -\lambda & 0 & 0 \\ 0 & 2-\lambda & 6 \\ 0 & 0 & 4-\lambda \end{pmatrix} = 0$$

$$= -\lambda(2-\lambda)(4-\lambda) = 0$$

$$\Rightarrow \lambda = 0, 2, 4$$

This is called the characteristic (polynomial) equation.

3) Solve $(A - \lambda I)\vec{x} = \vec{0}$

$$\lambda = 0$$

$$\begin{bmatrix} 0 & 0 & 0 & | & 0 \\ 0 & 2 & 6 & | & 0 \\ 0 & 0 & 4 & | & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 0 & 0 & 0 & | & 0 \\ 0 & 1 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{bmatrix}$$

$$\begin{cases} x_1 = x_1, \text{ free} \\ x_2 = 0 \\ x_3 = 0 \end{cases}$$

$$\vec{x} \in \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\} \\ = \text{Span} \{1\}$$

$$\lambda = 2$$

$$\begin{bmatrix} -2 & 0 & 0 & | & 0 \\ 0 & 0 & 6 & | & 0 \\ 0 & 0 & 2 & | & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 0 & 0 & | & 0 \\ 0 & 0 & 1 & | & 0 \end{bmatrix}$$

$$\begin{cases} x_1 = 0 \\ x_2 = x_3 \\ x_3 = 0 \end{cases}$$

$$\vec{x} \in \text{Span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \right\} \\ = \text{Span} \{x\}$$

$$\lambda = 4$$

$$\begin{bmatrix} -4 & 0 & 0 & | & 0 \\ 0 & -2 & 6 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & | & 0 \\ 0 & 1 & -3 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

$$\begin{cases} x_1 = 0 \\ x_2 = 3x_3 \\ x_3 = x_3 \end{cases}$$

$$\vec{x} \in \text{Span} \left\{ \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix} \right\} \\ = \text{Span} \{3x + x^2\}$$